



# Cytokine Storm Syndrome in COVID-19 Patients: Characteristics and Diagnosis

Harfouch RM\*

Department of microbiology and biochemistry, Faculty of pharmacy, Al Sham private university, Latakia, Syria

\*Corresponding author: Harfouch RM, Department of microbiology and biochemistry, Faculty of pharmacy, Al Sham private university, Latakia, Syria; E-mail: [r.h.foph.lat@aspu.edu.sy](mailto:r.h.foph.lat@aspu.edu.sy)

## Abstract

Cytokine storm syndrome (CSS) is a serious condition induced by a cascade of cytokine activation, characterized by general systemic inflammation, hyperferritinaemia, haemodynamic instability and multiple organ failure (MOF). Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged in Wuhan China, at the end of 2019, and rapidly developed into a global pandemic. This infection causes a dramatic increase of inflammatory cytokines in patients, suggesting the development of cytokine storm in some critical illness patients. Here, we summarize the pathogenesis, clinical manifestation of CSS, and highlight the current knowledge about the recognition and diagnosis of CSS in COVID-19.

**Keywords:** Cytokine storm syndrome (CSS); Coronavirus SARS-CoV-2; Characteristics

## Introduction

The coronavirus SARS-CoV-2 has infected more than 250 million people causing death to over than 5 million, with a mortality rate of approximately 2% worldwide. Many patients who die from COVID-19 suffer from hyper-inflammation with features of cytokine storm syndrome (CSS) and associated acute respiratory distress syndrome [1]. Only anti-inflammatory treatment have improved survival in these patients, although the antiviral Remdesivir was shown to reduce the length of hospital stay for COVID-19 patients [2]. The greatest survival benefit has been found with immunosuppressive glucocorticoids when given to those with an oxygen requirement. However, in the absence of an oxygen requirement or substantial systemic inflammation, patients treated with glucocorticoids may fare worse than those who receive standard care. For survival benefit, the selection of patients and timing of the administration of glucocorticoids is critical. Optimal treatment is earlier use of targeted anti-cytokine therapy to prevent CSS, without increasing viral replication [3].

## Definition of Cytokine storm syndrome (CSS)

Cytokine storm syndrome (CSS) is characterized by immune dysregulation due to continuous activation of lymphocytes and

macrophages, secretion of large amounts of cytokines including IL-1 $\alpha$ , IL-1 $\beta$ , IL-6, IL-18 and TNF- $\alpha$  and finally, overwhelming systemic inflammation and multi-organ failure with high mortality [4]. The term CSS was first used to describe the hypercytokinemia in graft after allogeneic stem cell transplant. Many viral, bacterial and parasitic infections can cause CSS such as Epstein-Barr virus (EBV) and Mycobacterium tuberculosis, which cause pathological immune activation characterized by elevated cytokines mainly interferon- $\gamma$  (IFN- $\gamma$ ) and soluble interleukin-2 receptor (sIL-2r) in patients with immune defects [5]. Levels of inflammatory cytokines IL-6, IL-8, MCP-1 and TNF $\alpha$  are significantly increased in COVID-19 plasma. Detailed correlation analysis revealed strong associations between some of the endothelial derived markers but only weak associations with age or general inflammatory parameters such as CRP. Fibrinogen levels correlate with increased P-selectin. In addition, plasma levels of IL-8 were strongly associated with circulating vWF levels. Local as well as systemic, circulating inflammatory markers coincide with various markers revealing endothelial activation and damage in COVID-19 patients.

## Pathogenesis of CSS

**Received date:** 07 December 2021; **Accepted date:** 21 December 2021; **Published date:** 28 December 2021

**Citation:** Harfouch RM (2021). Cytokine Storm Syndrome in COVID-19 Patients: Characteristics and Diagnosis. SunText Rev Biotechnol 2(2): 131.

**DOI:** <https://doi.org/10.51737/2766-5097.2021.031>

**Copyright:** © 2021 Harfouch RM. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The pathogenesis of CSS has not been fully known. Previous studies have shown that the development of CSS involves the imbalance of pro-inflammatory and anti-inflammatory mechanisms and the interaction of a variety of cells and cytokines, resulting in immune regulation disorder, causing a series of clinical manifestations. Previous studies have shown that a variety of cytokines can be markedly increased in patients with CSS, which may vary according to the heterogeneity of the disease background. In the setting of influenza virus-related CSS, a previous study found that patients infected with H5N1 had higher levels of monocyte chemoattractant protein 1 (MCP-1), interferon-gamma-induced protein-10 (IP-10) and IL-8 than patients infected with seasonal H1N1 influenza [6]. Moreover, previous studies also confirmed that cytokines played an important role in the pathogenesis of severe CoV infection. The serum proinflammatory cytokines (IFN- $\gamma$ , transforming growth factor- $\beta$  (TGF- $\beta$ ), IL-1, IL-6, IL-12) in severe SARS patients were significantly higher than those with mild to moderate symptoms, and serum proinflammatory cytokines (IFN- $\alpha$ , IL-6, IL-8) in severe MERS patients were significantly increased as well [7].

### Mechanisms of CSS in COVID-19

The cytokine storm in COVID-19 may have some differences from the cytokine storms in other clinical settings. Remarkably, the autopsy findings revealed that the lymphoid tissues and organs had been destroyed in COVID-19 patients, which is very unusual from CSS in sepsis and CAR T-cell therapy. Spleen atrophy and lymph node atrophy are observed in patients with COVID-19, whilst in other CSS-related diseases, lymphadenopathy and splenomegaly are more common. However, the specific mechanisms for these differences remain unclear and need to be further clarified [8]. Coronaviruses (CoVs) are enveloped, positive-sense, single-stranded RNA viruses, which have caused two large-scale pandemics in the last two decades, SARS and MERS [9]. Spike (S) proteins of coronaviruses, including the SARS-CoV, facilitate viral entry into their target cells via the interaction with functional cellular receptor identified as angiotensin-converting enzyme 2 (ACE2), which is highly expressed in alveolar epithelial cells, vascular endothelial cells, intestinal epithelial cells and renal proximal tubular cells. Functionally, ACE2, belonging to the ACE family, inactivates angiotensin II (Ang II) and generates angiotensin 1–7, a biologically active heptapeptide characterized by a potent vasodilator function. It has been demonstrated that the binding of the coronavirus spike protein to ACE2 leads to the down-regulation of ACE2, which in turn results in excessive production of vasoconstrictor Ang II and reduced production of vasodilator angiotensin 1–7.

Ang II also plays the role as a proinflammatory cytokine via angiotensin receptor 1 (AT1R). The Ang II-AT1R axis further

activates NF- $\kappa$ B and metalloprotease 17 (ADAM17), which stimulates the production of the mature form of epidermal growth factor receptor (EGFR) ligands and TNF- $\alpha$ . The activation of both NF- $\kappa$ B and STAT3, which in turn activate the IL-6 amplifier (IL-6 Amp), a mechanism for the hyper activation of NF- $\kappa$ B by STAT3, will lead to a hyper inflammatory state, resulting in increased pulmonary vascular permeability [10]. A retrospective study also found higher plasma concentrations of IL-2, IL-7, IL-10, IP-10, MCP-1, MIP-1a and TNF- $\alpha$  in intensive care unit (ICU) patients compared with no severe patients, suggesting there might be a cytokine storm in the body of severe patients [11].

### Diagnosis of CSS in COVID-19

There is no standard for the diagnosis of COVID-19 associated with CSS, and further clinical and laboratory investigations are needed. Here is a basic principle for consideration of CSS in COVID-19:

- A sudden or rapid progression with multiple organ involvement (such as liver, cardiac or renal injury)
- A significant decline of peripheral blood lymphocyte counts
- The significant elevation of systematic inflammatory indicators (such as CRP, serum ferritin, erythrocyte sedimentation rate)
- The elevation of multiple cytokines, such as IL-1 $\beta$ , IL-2R, IL-6, IFN- $\gamma$ , IP-10, MCP-1 and TNF- $\alpha$ .

Clinicians should keep highly alert on the possibility of CSS under these circumstances. However, given that CSS is a highly heterogeneous disease and may present with unspecific syndromes, the diagnosis of CSS in COVID-19 is very challenging and the development of a specific diagnostic test that helps to make the diagnosis of CSS earlier is a high priority for future research [12]. The inflammatory disorders in COVID-19 have been reported in many clinical studies. The COVID-19 causes a decrease of lymphocyte count and an increase of C reactive protein (CRP), especially in severely ill patients. The major subsets of the T lymphocytes (T cell) (CD3+ CD4+ T cell and CD3+ CD8+ T cells) are reduced in the COVID-19 and are significantly lower in the severe cases. The results of the other immune cells, the B cell and natural killer (NK) cell, have more inconsistency in recent researches. IL-6 was observed increased [13].

### Conclusion

The clinical, immunological, and pathologic features of COVID-19 have something in common with SARS and MERS. All the viruses can cause lymphopenia and influenza-like symptoms in the early stage. SARS and COVID-19 do not lead to the elevation of TNF- $\alpha$ , but the increase of IL-6 and IL-10 is more common in COVID-19. The IL-6 plays a crucial role in the pathologic of COVID-19, including the chemotaxis of neutrophils and

lymphocyte necrosis. Importantly, COVID-19 is more able to cause cytotoxic lymphocytes exhaustion.

## References

1. Harfouch RM, Moualla YM, Shahoud SM, Belal LA, Mohammad BS, Mustafa NM, et al. Epidemiology of COVID-19 in the most pandemic Countries: A review article. *Acad J Biotechnol.* 2021; 9: 21-27.
2. Harfouch RM, Alshaikh S, Alshimaly M, Assaad A, Ahmad J, et al. Therapeutic approaches for COVID 19: Challenges and successes. *Ann Clin Anal Med.* 2020.
3. Cron R, Caricchio R, Chatham WW. Calming the cytokine storm in COVID-19. *Nat Med.* 2021; 27: 1674-1675.
4. Shimizu M. Clinical features of cytokine storm syndrome. In: Cron RQ, Behrens EM, eds. *Cytokine Storm Syndrome.* Cham, Springer Int Publishing. 2019; 31-41.
5. Chen LYC, Hoiland RL, Stukas S, Wellington CL, Sekhon MS. Confronting the controversy: interleukin-6 and the COVID-19 cytokine storm syndrome. *Eur Respir J.* 2020; 56.
6. Jong MD, Simmons CP, Thanh TT, Hien VM, Smith GJDF, Chau TNB, et al. Fatal outcome of human influenza A (H5N1) is associated with high viral load and hypercytokinemia. *Nat Med.* 2006; 12: 1203-1207.
7. Channappanavar R, Perlman S. Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. *Semin Immunopathol.* 2017; 39: 529-539.
8. Yao XH, Li TY et al. A pathological report of three COVID-19 cases by minimally invasive autopsies. *Chin J Pathol.* 2020; 49: 9.
9. Alhoury A, Salloum A, Harfouch RM, Soumya G. Possible side effects of using detergents during the Covid19 pandemic in Syria. *Ann Clin Cases.* 2020; 1: 1023.
10. Murakami M, Kamimura D, Hirano T. Pleiotropy and specificity: insights from the interleukin 6 family of cytokines. *Immunity.* 2019; 50: 812-831.
11. Huang C, Wang Y, Li X et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020; 395: 497-506.
12. Gao YM, Xu G, Wang B, Liu BC. Cytokine storm syndrome in coronavirus disease 2019: a narrative review. *J Intern Med.* 2020; 289: 147-161.
13. Tang Y, Liu J, Zhang D, Xu Z, Ji J, et al. Cytokine Storm in COVID-19: The current evidence and treatment strategies. *Front Immunol.* 2020; 11: 1708.